

STOP OVERDOSE

RECOGNIZE THE SIGNS OF OPIOID OVERDOSE

An opioid overdose
requires immediate
medical attention.



Face is clammy to touch and has lost color. Vomiting or gurgling noises.



Body is limp. Fingernails or lips have a blue or purple tinge.



Sleep is deep and cannot be awoken. Difficulty speaking.



Breathing is slow or has stopped.



Heartbeat is slow or has stopped.

DON'T RUN. CALL 911. YOU CAN SAVE A LIFE.

The ND Good Samaritan Law protects you so you can protect your friend.

20

DEATHS
IN 2013

61

DEATHS
IN 2015

**Overdose deaths in North Dakota increased
from 20 deaths in 2013 to 61 deaths in 2015.**

CDC/NCHS, National Vital Statistics System, Mortality

Source: Substance Abuse and Mental Health Services Administration. SAMHSA Opioid Overdose Prevention Toolkit. HHS Publication No. (SMA) 16-4742. Rockville, MD: Substance Abuse and Mental Health Services Administration, 2016.



For more information, visit
prevention.nd.gov/stopoverdose

Created in partnership with the Reducing Pharmaceutical
Narcotics in our Communities Task Force

NORTH DAKOTA
PREVENTION
RESOURCE & MEDIA CENTER